

TCIメール

2018.7 **178**



▼ 目次


2 寄稿論文

- Carbocyclization of Carbohydrates to Hydroxylated Cycloalka(e)nes

Tony K. M. Shing
Professor, Department of Chemistry,
Faculty of Science and Technology, Keio University

28 製品紹介

- 筒状炭素分子:(6,6)カーボンナノベルト
- 高性能な液晶性半導体材料:Ph-BTBT-10
- 光学活性なシクロプロパンの合成に有用な不斉ルテニウム触媒
- ペルオキシダーゼ用調製済み基質溶液
- c-Jun N末端キナーゼ(JNK)阻害剤:SU 3327
- シクロオキシゲナーゼ(COX)阻害剤:フルルビプロフェン



Carbocyclization of Carbohydrates to Hydroxylated Cycloalka(e)nes

Tony K. M. Shing

*Department of Chemistry, Faculty of Science and Technology, Keio University**Email: shing@chem.keio.ac.jp*


Abstract: Carbohydrates, inexpensive and rich in stereochemistry, are nature's gifts to the synthetic organic chemists. We have been conducting research on the carbocyclization of carbohydrates for many years, i.e. the conversion of simple, readily available monosaccharides into hydroxylated cycloalkanes and cycloalkenes with pharmaceutical potential. Several key reactions are presented in this Article to illustrate such facile transformation namely $[\pi 4s+\pi 2s]$ cycloaddition - intramolecular nitron-alkene and nitrile oxide-alkene cycloadditions, intramolecular direct aldol reaction and intramolecular Horner–Wadsworth–Emmons olefination. These protocols provide facile entries to 6 and 7-membered hydroxylated carbocycles in enantiomerically pure forms which then could be further elaborated into target molecules. Examples including calystegines, tropane alkaloid, cyclohexanyl and cyclohexenyl gabosines, cycloheptanones, valiolamine, validoxylamine G, pseudo-acarviosin, and a SGLT2 inhibitor are presented.

Keywords: carbohydrates, natural products, organic synthesis, pericyclic reactions, hydroxylated carbocycles

This review article describes our research efforts on finding a short, facile, and efficient way of transforming carbohydrates into hydroxylated carbocycles over a period of about 30 years. There are excellent reviews on the topic published elsewhere.¹


The story begins when I was a postdoctoral fellow at Oxford working with Professor George Fleet in 1983. We were trying to synthesize shikimic acid from D-mannose using an intramolecular aldol condensation or a Wittig-type alkenation as the key step. Installing a phosphono-acetate moiety at the C-5 of the benzyl furanoside intermediate **1** to form the Wittig–Horner–Emmons precursor met with difficulty (Scheme 1).²

Simple S_N2 displacement reactions in carbohydrates are known to be sluggish/difficult due to the high concentration of oxygen functions (electron rich). Thus the derived standard electrophiles, 5-tosylate **2** and 5-iodide **3** were found to be inert towards substitution. We then attempted to use the 5-triflate **4** as a reactive leaving group (this was new at that time). Gratifyingly, the alkylation proceeded smoothly to give the Horner–Wadsworth–Emmons (HWE) alkenation precursor in a good yield. Subsequent unmasking the aldehyde function in **5** by hydrogenolysis afforded lactol **6** that was treated with base to give the cyclohexene motif, the protected shikimic acid **7**. Acid catalysed removal of the blocking groups in **7** furnished shikimic acid in an excellent overall yield.²




Up to date, this intramolecular HWE alkenation is still one of the most powerful strategies for the carbocyclization of sugar molecules. A notable example of using this strategy was demonstrated by Fang and co-workers³ to synthesize anti-viral agent Tamiflu[®] **12** and its more potent phosphonate analog **13** in a similar manner (Scheme 2).

* Tamiflu[®] is a registered trademark of F. Hoffmann-La Roche, Ltd.




The reasons for researching on carbocyclization of carbohydrates are attributed to the fact that many biologically active molecules are hydroxylated cycloalkanes/alkenes. Carbohydrates are obvious starting materials that offer the most facile and economical way to approach these highly oxygenated carbocycles with defined stereochemistry.


Some hydroxylated cyclohexanes and cyclohexene are listed in Figure 1. They display a wide plethora of bioactivities including antibiotic, antitumor, antiviral, glycosidase inhibitory activities.



Calystegines,⁴ displayed specific glycosidase inhibition, are a group of bicyclic heterocycles which are in equilibrium with the open chain form, a hydroxylated gamma-amino cycloheptanone that are potentially accessible from carbohydrates (Figure 2).



Gabosines are a group of hydroxylated cyclohexanones/cyclohexenones that display antibiotic, anticancer, and DNA binding properties (Figure 3).⁵ All these compounds should in theory be approachable from sugars, but how could we effect the carbocyclization efficiently?




Towards that end, we have been investigating the following strategies for the carbocyclization of sugars:

1. Carbocyclization of Carbohydrates *via* an Intramolecular Nitrone-Alkene Cycloaddition (INAC)
2. Carbocyclization of Carbohydrates *via* an Intramolecular Nitrile Oxide-Alkene Cycloaddition (INOAC)
3. Carbocyclization of Carbohydrates *via* an Intramolecular Direct Aldol Addition of Sugar Diketone
4. Carbocyclization of Carbohydrates *via* an Intramolecular Horner–Wadsworth–Emmons (HWE) Olefination


From 1984, I started my independent research career at the University of Manchester and began to investigate the construction of carbocycles using an INAC reaction as the key step.

1 Carbocyclization of Carbohydrates *via* an Intramolecular Nitrone-Alkene Cycloaddition (INAC)

2,3-*O*-Isopropylidene D-ribose **23** underwent a chelation controlled Grignard vinylation to give triol **24** (Scheme 3). Glycol cleavage oxidation of the vicinal diol in **24** afforded lactol **25** which on heating with *N*-methyl hydroxylamine furnished a hydroxylated cyclopentane **27** smoothly. This is an extremely short approach towards 5-membered carbocycles and its application to the syntheses of carbocyclic nucleosides should have received more attention.⁶




On the other hand, chelation controlled vinylation of diacetone mannose **28** gave allyl alcohol **29** stereoselectively which was protected as benzyl ether **30** (Scheme 4). Regioselective mild acid hydrolysis of the terminal acetonide was feasible as the *O*-8 was the least hindered and therefore protonated first. The resultant diol **31** was oxidatively cleaved to afford aldehyde **32** which was reacted with *N*-methyl hydroxylamine to give nitron **33**. INAC occurred smoothly on heating to give cyclohexane **34** in good yield. In both cases, the new C–N bond is *anti* to the *O*-2 stereochemistry and the ring fusion is *cis*.⁶




We had more interests in this INAC reaction because we reasoned that it could be controlled to provide 7-membered carbocycles. When a sugar molecule was elaborated to have a terminal alkene, on reaction with *N*-alkyl hydroxylamine, there would be two possible modes of INAC cyclization, the *exo* or the *endo* mode (Scheme 5), which leads to either a fused or a bridged isoxazolidine, respectively.⁷ To be specific, a hept-6-

enose would provide either a fused bicyclo[4.3.0] system, i.e. a cyclohexane skeleton from an *exo*-mode or a bridged bicyclo[4.2.1] system, i.e. a cycloheptane skeleton from an *endo*-mode INAC cyclization. We were particularly intrigued by the potential formation of 7-membered carbocycles as there are few synthetic strategies towards cycloheptanes whereas the synthetic methods for the construction of 5- and 6-membered carbocycles are plentiful.




We realised that the formation of a cyclohexane skeleton *via* the *exo*-mode of cyclization of hept-6-enose is the usual regioselectivity outcome with conventional protecting groups like benzyl ether, silyl ether, acetonide, and esters *etc.* For examples, nitronium **35** and **36**, both with a *cis*-diol acetonide protecting group gave cyclohexane rings in high yields (Scheme 6).⁸ The stereochemistry of 4-OH apparently had no effect on the regioselectivity.




We reasoned that the mode of cyclization might be controlled by the blocking groups as they affect the orbital overlap between the nitronium and the alkene. Thus we investigated the effect of *trans*-diacetal blocking group on the regioselectivity of hept-6-enose. Epimeric nitronium **37** and **38** were readily prepared from D-arabinose involving glycosidation with benzyl alcohol, diacetalisation, debenzoylation, Grignard allylation, glycol cleavage oxidation and reaction with *N*-methyl hydroxylamine. Interestingly, nitronium **37** and **38** with a *trans*-diacetal blocking group (the diol moiety is *trans*) afforded *endo*-mode of cyclization products **41**, **44**, **45**, i.e., cycloheptane rings, for the first time (Scheme 7). Another notable feature of this reaction is that a *trans*-fused


bicyclo[4.3.0] system, i.e. cyclohexane **40** and **43** were formed. The lower yield of *cis*-fused **42** and higher yield of *trans*-fused **43** is probably attributed to the 1,3 diaxial interaction between 4-OH and the 6-hydroxymethyl in **42**, this steric interaction is absent in the *cis*-cycloadduct **39**. The rigidity of the diacetal ring requires 2-O and 3-O in the *trans*-diequatorial dispositions, thus 4-OH and the 6-hydroxymethyl in **42** must be axial.⁸




We reasoned that cycloheptane ring might be more accommodating than 6-membered rings as far as ring strain is concerned. We therefore went on to study the effect of an acetone with a *trans*-diol blocking group on the regioselectivity of nitron cycloaddition of hept-6-enose (Scheme 8). Nitrones **46** and **47** with an acetone of *trans*-2,3-diol provided *endo*-cycloadducts **48** and **49** as the major products for the first time.⁹




Hept-6-enose nitrones **50** and **51** with a 3,4-*trans*-acetonide were even more remarkable as only the *endo*-mode cycloheptanes were harvested in good yields (Scheme 9).¹⁰




We were wondering if the stereochemistry of the C-2/C-5 hydroxyl groups might affect the regioselectivity of the INAC reactions. We thus constructed a hept-6-enose nitronium **60** with only a 3,4-*trans*-acetonide from L-tartaric acid using standard reactions (Scheme 10). Only a cycloheptane ring **61** was obtained, thereby confirming that the *endo*-mode of cyclization was controlled by the *trans*-acetonide ring.¹¹



Since we have established that 7-membered cycloheptanes can be accessed *via* a *trans*-acetonide directed *endo*-selective INAC reactions of hept-6-enoses, we set out to synthesize calystegine analogs. In this way, D-xylose was transformed into cycloheptanes **63**, **64**, **65** with a 3,4-*trans*-acetonide (Scheme 11). Debenzoylation afforded the corresponding alcohols in which **66** and **68** were oxidized to the same ketone **69**. Acidic deacetonation of **69** followed by global hydrogenolysis furnished (*S*)-3-hydroxy-calystegine B₅ **71**.¹⁰ We found that *tert*-butanol was a good solvent for hydrogenolysis of a *N*-benzyl group as no side products from *N*-alkylation was possible.




On the other hand, D-ribose was converted into aldehyde **72** which underwent *trans*-acetone directed *endo*-selective INAC reaction to give cycloheptane **73** in an excellent yield (Scheme 12). Acid hydrolysis of the acetonide ring produced the diol **74** which underwent steric controlled regio-selective oxidation of the less hindered alcohol to give ketone **75**. Hydrogenolysis of the N–O bond afforded a B-type calystegine analog **76**.¹⁰




Then we investigated this *endo*-selective INAC approach towards syntheses of tropane alkaloid **90** and analogs.

(A) Strategy towards tropane alkaloid **1**: INAC of a Nitron with a 2,3-*cis*-Isopropylidene as Blocking Group and an α,β -Unsaturated Ester as Dipolarophile




Since the alkene moiety in **77** is an electrophilic alkene whereas the alkenes used in our INAC cyclization mode studies were simple nucleophilic alkenes, we were hoping that the electrophilic alkene of the enoate group in **77** might cause an *endo*-selective INAC reaction attributable to HOMO–LUMO reasons (Scheme 13). This proposal was supported by a precedent¹² that in the intermolecular nitron-alkene cycloaddition of α,β -unsaturated ester **82**, and nitron **81**, the new C–O was installed on the β -carbon in cycloadducts **83** and in **84** (Scheme 14). If this were applied to our INAC case, the *endo*-mode cyclization would have taken place. We were excited with this precedent and therefore proceeded with the synthesis.



Scheme 14. Intermolecular cycloaddition of nitron **81** and enoate **82**


To our disappointment, the INAC of nitron **85** produced only *exo*-mode cycloadducts **86** and **87**, 6-membered carbocycles. Hence it appears that steric control is more important than electronic control for the regioselectivity (*exo* versus *endo*) of INAC reactions of hept-6-enes (Scheme 15).¹¹



Scheme 15. INAC of enoate with a *cis*-acetonide


(B) Strategy towards tropane alkaloid 2: INAC of a Nitron with a 3,4-*trans*-Isopropylidene as Blocking Group and an α,β -Unsaturated Ester as Dipolarophile

Since we had shown that 3,4-*trans*-acetonide would direct an *endo*-selective mode of cyclization, we devised a synthetic route towards cocaine based on this key step as summarised in Scheme 16.




Scheme 16. Steric controlled synthetic plan towards tropane alkaloid **90**

Capitalising from our studies that 3,4-*trans*-acetonide would produce cycloheptane ring on INAC reactions of hept-6-enoses, we proceeded to prepare the nitron **97** from ribose.¹³ The known allyl alcohol **91** was readily obtained from D-ribose *via* a steric-controlled allylation (Scheme 17). Thermodynamic acetonation of **91** formed the 7,8-*O*-isopropylidene first as the primary 8-alcohol was the most reactive of all. Then an acetonide was formed between *O*-4,5 instead of *O*-5,6 because the substituents on the acetonide ring **92** were *trans* to each other. The diacetonide **92** was then transformed into diol **95** without incident. Glycol cleavage oxidation of **95** with silica supported NaIO₄ furnished aldehyde **96** smoothly. This solid-phase NaIO₄ reagent¹⁴ is highly recommended as the operation is very simple and the aldehyde harvested is usually un-hydrated. INAC reaction of **97** did produce the desired cycloheptane derivative **98** as the major product.¹³




Scheme 17. INAC synthetic route towards tropane alkaloid

As the cycloheptane skeleton was installed correctly, cycloadduct **98** was converted into natural tropane alkaloid **90** and its analogs **100–104** as shown in Figure 4.¹³




2 Carbocyclization of Carbohydrates *via* an Intramolecular Nitrile Oxide-Alkene Cycloaddition (INOAC)


The use of chloramine-T in the conversion of ene-sugar oxime into a nitrile oxide and its subsequent INOAC reaction is known.¹⁵ This cycloaddition shown in Scheme 18 can only be *exo*-mode as the *endo*-mode of cyclization would produce an impossible C=N bond at the bridge head position (Bredt's rule).¹⁶



Common chemicals for the transformation of oxime into nitrile oxide include NaOCl, NaOCl/NEt₃ and *N*-chlorosuccinimide/NEt₃.¹⁶ Under the basic conditions, the free hydroxyl group in nitrile oxide **105** would attack the nitrile carbon to give oximolactone **106** and failed to produce a carbocycle. The hydroxy groups of the sugar have to be protected in order to achieve satisfactory yields. Thus, providing a mildly acidic environment for the INOAC reactions is crucial in achieving a fruitful carbocyclization without the formation of oximolactone.




We found that adding flash chromatography silica gel to “buffer” the reaction medium dramatically increased the yield of the chloramine-T mediated INOAC reaction from 62 to 94% (Scheme 20).¹⁷




Towards the construction of 5-membered carbocycles, similar improvement in reaction yields were observed and examples are shown in Scheme 21. It is interesting to note that other acidic conditions attempted including the addition of acetic acid, benzoic acid, acetate buffer or phosphate buffer did not cause the INOAC to occur.


The silica gel mediated INOAC reactions were compatible with substrates containing unprotected hydroxyl groups, thus reducing masking/unmasking steps and rendering a synthesis shorter.¹⁷



We have applied this methodology to the construction of gabosine O and F.¹⁸ Mannose was transformed readily into oxime **114** with a free hydroxyl group (Scheme 22). INOAC of **114** occurred smoothly to give a 6-membered carbocycle **115** which was hydrogenolysed, dehydrated, and then hydrogenated from the less hindered side to form ketone **118**. Acidic hydrolysis of the acetone in **118** yielded the target molecule gabosine O.¹⁸




On the other hand, L-arabinose was transformed into oxime **119** with a free hydroxyl group. INOAC reaction of **120** furnished the 6-membered carbocycle in an excellent yield with the new C–C bond equatorially installed. Standard conversion afforded the target molecule gabosine F (Scheme 23).



3 Carbocyclization of Carbohydrates *via* an Intramolecular Direct Aldol Addition of Sugar Diketone

First of all, we started our investigation on D-glucose since voglibose **18** and valiolumine **19** are obvious synthetic targets. Glycosidation of D-glucose with allyl alcohol and then diacetalization produced 2,3- **124** and 3,4-diacetals **123** which were readily separable upon acetonation to the 4,6-acetonide **126**.¹⁹ Palladium catalysed deallylation of glycoside **126** gave lactol **127** which underwent Grignard methyl addition followed by PDC oxidation afforded the 2,6-diketone **129**. Intramolecular aldol addition reactions of **129** were carried out under various basic conditions and the best results are shown in Table 1.



The stereo-selective formation of a particular aldol product can be controlled by either L-proline, strong or weak amine base. It is noteworthy that all the direct aldol reactions are not reversible. The resistance towards a retro-aldol reaction is probably attributable to steric reasons.





Table 1. Summary of direct aldolization conditions of 2,6-diketone **129**


Entry	Conditions	Results		
		130	131	132
1	L-Proline (0.3 eq), DMSO	82%	8%	2%
2	KHMDS (1 eq), Toluene, -78 °C	–	75%	–
3	Et ₃ N (1.5 eq), CH ₂ Cl ₂	–	–	95%

The preparation of mannose diketone **135** was executed in a similar manner (Scheme 25). Disappointingly, all strong and weak basic conditions failed to generate an aldol product with the exception of L-proline, which provided aldol **136** as the sole cyclohexanone in 60% yield.¹⁹



Scheme 25. Preparation of aldol **136** from D-mannose


We have also investigated an intramolecular direct aldolization of 2,7-diketones, in search for a synthetic avenue towards hydroxylated cycloheptanones. The 2,7-diketones were prepared from D-ribose or D-mannose using standard reactions and the results are summarized in Scheme 26.²⁰



Application of the versatile intramolecular direct aldolization strategy to synthesis was demonstrated by the following examples.


3a. Synthesis of Valiolamine and Voglibose (AO128)

D-Glucose was transformed smoothly into aldol **131** as stated above. Oximation of the ketone moiety in **131** followed by hydrogenation/hydrogenolysis furnished amine **142**. Acidic removal of the acetals in **142** provided valiolamine **19** which, reportedly, could be converted into the anti-diabetic agent voglibose (AO128) **18** (Scheme 27).¹⁹




3b. Synthesis of Pseudo-Acarviosin


Methyl acarviosin **143** was obtained from the natural anti-diabetic agent, acarbose, by acidic methanolysis (Figure 5). As methyl acarviosin is a stronger α -amylase inhibitor, we wanted to replace the sugar moiety in **143** with a cyclohexane to render the compound more stability in acid conditions. In search for an improved anti-diabetic drug, we proceeded to synthesize pseudo-acarviosin **144** for biological screening.



Retrosynthesis of pseudo-acarviosin **144** *via* a palladium catalysed allylic substitution would give two fragments, an allylic chloride **145** and a cyclohexanyl amine **146**. Chloride **145** would be accessed from D-glucose *via* a direct aldol strategy while the amine **146** should be approached from L-arabinose through an INAC reaction.²¹




The palladium-catalysed coupling plan is the result of a massive experimentation. We attempted a great many coupling reaction conditions between different partners, amine with mesylate, ketone, allylic epoxide, allylic cyclic sulphite, or allylic acetate.²² As most of the electrophiles are unstable under the conditions and elimination is the major side-reaction, none of the coupling reactions gave the desired product except for using allylic chloride as the coupling partner shown in Scheme 29.²³




Scheme 29. Successful coupling reactions when allylic chloride was used

Since the coupling conditions were established, we started to construct the two coupling partners **145** and **146**. D-Glucose was converted into aldol **130** as said earlier.¹⁹ Elimination of the alcohol in **130** followed by reduction of the ketone group in **147** produced allylic alcohol **148** which was separated from its epimer *via* an acetylation and deacetylation protocol. The isolated **148** was then transformed into the allylic chloride **145** by standard reactions (Scheme 30).


Scheme 30. Preparation of allylic chloride **145** from D-glucose

On the other hand, the amine coupling partner **146** was assembled from L-arabinose *via* an INAC reaction of a protected hept-6-ene **151** as the key step (Scheme 31).²¹ The major *exo*-cycloadduct **154** with a *trans*-ring fusion was formed with the correct stereochemistry. It is noteworthy that the *trans*-fused *exo*-cycloadduct could only be harvested with the nitrone containing a *trans*-diacetal blocking group. The cycloadduct **154** was transformed readily into the cyclohexyl amine **146** using standard reactions with a good overall yield.



Scheme 31. Preparation of amine **146** from L-arabinose


With the coupling partners in our hand, palladium catalysed allylic substitution of chloride **145** with amine **146** proceeded with retention of configuration to give the protected allylic amine **161** (Scheme 32). Hydrolysis gave the target molecule pseudo-acarviosin **144** which was shown to be a stronger sucrose and glucoamylase inhibitor than acarbose.²¹



Scheme 32. Synthesis of pseudo-acarviosin **144**


3c. Synthesis of Validoxylamine G

Capitalising on the palladium-catalysed allylic substitution reaction, the silyl ether **162** of the afore-prepared amine **142** and allylic chloride **145** (both are direct aldol products from D-glucose) were coupled together to form pseudo-glycosyl amine **163** which on complete acidic deprotection afforded validoxylamine G **164** in a good yield (Scheme 33).¹⁹




3d. Synthesis of Gabosines

A number of gabosines were easily assembled from D-glucose *via* the direct aldol strategy. Only a few of them are illustrated here.^{24,25} Thus the aforesaid enone **147** was regio- and stereo-selectively reduced to the α -allylic alcohol **165** in an excellent yield (Scheme 34). Functional group manipulation of **165** with standard reactions then furnished gabosine A, D, and E without incident.²⁵




4 Carbocyclization of Carbohydrates via an Intramolecular Horner-Wadsworth-Emmons (HWE) Olefination

Intramolecular HWE olefination was introduced at the beginning of this article, interestingly, it is perhaps still the best strategy for a short and efficient carbocyclization of carbohydrate. Towards this end, we globally protected D-gluconolactone with 2-methoxypropene as mixed acetals **166** (Scheme 35).²⁶ Addition of methyl phosphonate carbanion followed by oxidation of the alcohol in **167** caused intramolecular HWE alkenation to proceed concomitantly to give the protected cyclohexanone **169** in just 3 steps from the free sugar! Acidic removal of the acetals afforded gabosine I **170** and regioselective acetylation at the primary alcohol provided gabosine G **171**. The relatively labile mixed acetal blocking groups were problematic, but we have circumvented this obstacle with ethoxymethyl (EOM) protecting group.²⁷



Since EOM protected cyclohexanone **172** could also be synthesized readily from D-gluconolactone (Scheme 36), application of this route to the syntheses of streptol and other gabosines were accomplished.^{27,28} Exploiting this facile carbocyclization strategy, we embarked on the construction of sodium-dependent glucose cotransporter 2 (SGLT2) inhibitors.



Sergliflozin is a potent SGLT2 inhibitor, but was abandoned for further investigation because phenyl glycosides are prone to hydrolysis and therefore not suitable for development into useful medicine. Replacing the sugar moiety in sergliflozin with a pseudo-sugar unit would provide a stable “glycoside” **173** as the “glycosidic bond” is now an ether linkage (Figure 6).²⁹





Figure 6. Structures of sergliflozin and pseudo-aryl glucoside **173**

The synthetic avenue towards pseudo glucopyranoside **173** is shown in Scheme 37. Regio- and stereo-selective reduction of **172** gave α -alcohol **174** which was activated and displaced with chloride to β -chloride **175**. Palladium catalysed allylic substitution of the allylic chloride **175** with phenolic alcohol **176** proceeded with retention of configuration to form allylic ether **177** in an excellent yield. Catalytic hydrogenation followed by acid removal of the EOM groups afforded the target molecule **173** which is a potent and selective SGLT2 inhibitor. Other analogs have also been made on extending this carbocyclization strategy.^{30,31}



Scheme 37. Synthesis of pseudo-aryl glucopyranoside **173**

Conclusion

In this Article, transformation of carbohydrates into highly oxygenated cycloalkanes and cycloalkenes has been described using four synthetic strategies, namely intramolecular nitrone-alkene and nitrile oxide-alkene cycloadditions, direct aldol addition, and Horner–Wadsworth–Emmons (HWE) olefination. The latter strategy probably offers the most facile and efficient avenue to the pseudo-D-glucopyranose motif to date. The carbocyclized intermediates could then be elaborated into various target molecules containing hydroxyl/amine functional groups of defined stereochemistry. Considering the low cost, availability in large quantities, and rich stereochemistry of sugar molecules, carbohydrates are perhaps the ideal starting materials for the construction of heavily oxygenated natural products or molecules with pharmaceutical implication. Research towards the syntheses of carbocyclic nucleosides should have received more attention.

Acknowledgments

The author wishes to thank the University of Manchester, the Chinese University of Hong Kong and the Hong Kong Research Grant Council for supporting these works. Sincere gratitude goes to his research collaborators and co-workers for their efforts and stimulating ideas, especially to Professor Tohru Yamada of Keio University for his unflinching support, encouragement, and friendship.

References

- 1) (a) O. Arjona, A. M. Gómez, J. C. López, J. Plumet, *Chem. Rev.* **2007**, *107*, 1919. (b) R. J. Ferrier, S. Middleton, *Chem. Rev.* **1993**, *93*, 2779.
2. (a) G. W. J. Fleet, T. K. M. Shing, *J. Chem. Soc., Chem. Commun.* **1983**, 849. (b) G. W. J. Fleet, T. K. M. Shing, S. W. Warr, *J. Chem. Soc., Perkin Trans. 1* **1984**, 905.
3. J.-J. Shie, J.-M. Fang, S.-Y. Wang, K.-C. Tsai, Y.-S. E. Cheng, A.-S. Yang, S.-C. Hsiao, C.-Y. Su, C.-H. Wong, *J. Am. Chem. Soc.* **2007**, *129*, 11892.
4. (a) B. Dräger, *Nat. Prod. Rep.* **2004**, *21*, 211. (b) N. Asano, *Curr. Top. Med. Chem.* **2003**, *3*, 471.
5. (a) K. Tatsuta, T. Tsuchiya, N. Mikami, S. Umezawa, H. Umezawa, H. Naganawa, *J. Antibiotics* **1974**, *27*, 579. (b) G. Bach, S. Breiding-Mack, S. Grabley, P. Hammann, K. Hütter, R. Thiericke, H. Uhr, J. Wink, A. Zeeck, *Liebigs Ann. Chem.* **1993**, *3*, 241. (c) Y.-Q. Tang, C. Maul, R. Höfs, I. Sattler, S. Grabley, X.-Z. Feng, A. Zeeck, R. Thiericke, *Eur. J. Org. Chem.* **2000**, *1*, 149.
6. T. K. M. Shing, D. A. Elsley, J. G. Gillhouley, *J. Chem. Soc., Chem. Commun.* **1989**, 1280.
7. J. J. Tufariello, In *1,3-Dipolar Cycloaddition Chemistry*, ed. by A. Padwa, Wiley, New York, **1984**, Vol. 2, pp. 83-168.
8. T. K. M. Shing, A. W. F. Wong, T. Ikeno, T. Yamada, *J. Org. Chem.* **2006**, *71*, 3253.
9. T. K. M. Shing, A. W. F. Wong, T. Ikeno, T. Yamada, *Chem. Eur. J.* **2009**, *15*, 2693.
10. T. K. M. Shing, A. W. F. Wong, T. Ikeno, T. Yamada, *Org. Lett.* **2007**, *9*, 207.
11. T. K. M. Shing, K. H. So, unpublished results.
12. E. G. Baggolini, J. A. Iacobelli, B. M. Hennesy, A. D. Batcho, J. F. Sereno, M. R. Uskokovic, *J. Org. Chem.* **1986**, *51*, 3098.
13. T. K. M. Shing, K. H. So, *Org. Lett.* **2011**, *13*, 2916.
14. Y.-L. Zhong, T. K. M. Shing, *J. Org. Chem.* **1997**, *62*, 2622.
15. (a) A. Pal, A. Bhattacharjee, A. Bhattacharjya, A. Patra, *Tetrahedron* **1999**, *55*, 4123. (b) S. Majumdar, R. Mukhopadhyay, A. Bhattacharjya, *Tetrahedron* **2000**, *56*, 8945.

16. P. Caramella, P. Grünanger, In *1,3-Dipolar Cycloaddition Chemistry*, ed. by A. Padwa, Wiley, New York, **1984**, Vol. 1, pp. 291-392.
17. T. K. M. Shing, W. F. Wong, H. M. Cheng, W. S. Kwok, K. H. So, *Org. Lett.* **2007**, *9*, 753.
18. T. K. M. Shing, K. H. So, W. S. Kwok, *Org. Lett.* **2009**, *11*, 5070.
19. T. K. M. Shing, H. M. Cheng, *Org. Lett.* **2008**, *10*, 4137.
20. T. K. M. Shing, H. M. Cheng, *Org. Biomol. Chem.* **2015**, *13*, 4795.
21. T. K. M. Shing, H. M. Cheng, W. F. Wong, C. S. K. Kwong, J. Li, C. B. S. Lau, P. S. Leung, C. H. K. Cheng, *Org. Lett.* **2008**, *10*, 3145.
22. T. K. M. Shing, S. H.-L. Kok, unpublished results.
23. (a) T. K. M. Shing, S. H.-L. Kok, *Tetrahedron Lett.* **2000**, *41*, 6865. (b) T. K. M. Shing, C. S. K. Kwong, A. W. C. Cheung, S. H.-L. Kok, Z. Yu, J. Li, C. H. K. Cheng, *J. Am. Chem. Soc.* **2004**, *126*, 15990.
24. T. K. M. Shing, H. M. Cheng, *Synlett* **2010**, *21*, 142.
25. T. K. M. Shing, H. M. Cheng, *Org. Biomol. Chem.* **2009**, *7*, 5098.
26. T. K. M. Shing, H. M. Cheng, *J. Org. Chem.* **2007**, *72*, 6610.
27. T. K. M. Shing, Y. Chen, W.-L. Ng, *Synlett* **2011**, *22*, 1318.
28. T. K. M. Shing, Y. Chen, H. T. Wu, *Synlett* **2012**, *23*, 1793.
29. T. K. M. Shing, W.-L. Ng, J. Y.-W. Chan, C. B.-S. Lau, *Angew. Chem. Int. Ed.* **2013**, *52*, 8401.
30. W.-L. Ng, K.-M. Lau, C. B.-S. Lau, T. K. M. Shing, *Angew. Chem. Int. Ed.* **2016**, *55*, 13818.
31. W.-L. Ng, H.-C. Li, K.-M. Lau, A. K. N. Chan, C. B.-S. Lau, T. K. M. Shing, *Sci. Rep.* **2017**, *7*, 5581.

執筆者紹介

Tony K. M. Shing

Professor, D.Sc.

Department of Chemistry, Faculty of Science and Technology, Keio University



Professor Tony K. M. Shing obtained his first degree from the University of Hong Kong, his M.Sc. (analytical chemistry), Ph.D. (carbohydrate chemistry) and D.Sc. (organic synthesis) degrees from the University of London. After postdoctoral work at McGill University and Oxford University, Prof. Shing began his independent career at the University of Manchester England, as a New Blood Lecturer in 1984. He was visiting Associate Professor at the University of California, Irvine, before returning to Hong Kong in 1990. He joined the Department of Chemistry at the Chinese University of Hong Kong (CUHK) as a Lecturer, was promoted to Senior Lecturer in 1993, and to full Professor in 1996. In 2016 August, he retired from CUHK and joined Sophia University

as an Invited Visiting Professor. Professor Shing moved to Keio University in October last year and is now a full Professor at the Department of Chemistry. He is also the Chairman of the Hong Kong Section of the Royal Society of Chemistry. His research interest is on the syntheses of organic molecules with chemotherapeutic potential, namely anti-cancer, anti-viral, or anti-diabetic activity.

筒状炭素分子：(6,6)カーボンナノベルト

I1078 (6,6)Carbon Nanobelt Bis(tetrahydrofuran) Adduct (1)

10mg 12,000 円

カーボンナノベルトは、ベンゼン環が縮環して筒状の構造をもつ炭素分子の総称として提唱され、その歴史は 1954 年に文献に記載されたことに始まります¹⁾。その後、さまざまな筒状炭素分子が提案され、多くの化学者がそれらの合成に挑戦しました²⁾。1991 年にカーボンナノチューブが発見されると³⁾、カーボンナノベルトがカーボンナノチューブの部分構造であることが明らかになり、ますます注目を集めるようになりました。一方、カーボンナノベルトは、ベンゼン環が筒状に曲がることで大きなひずみが生じるため、有効な合成法はこれまで報告されていませんでした。

伊丹、瀬川らのグループは、ひずみのない環状分子を筒状構造に変換する方法で、パラキシレンから 11 段階で **1** を合成することに成功しました⁴⁾。得られた **1** の基礎物性は、各種分光学的手法により解析されました。X 線を用いた構造解析法から、カーボンナノベルトはカーボンナノチューブと同様の筒状構造をもつことが明らかになりました。また、可視光の吸収および蛍光の分析からは、構造の剛直さや筒状構造全体に電子が通る性質が観測されるとともに、(6,6)カーボンナノチューブに非常に近い性質をもつ分子であることがラマン分光法によって実証されました⁴⁾。

カーボンナノベルトは、単一構造からなるカーボンナノチューブの合成の実現や、新しい機能性材料開発の道を拓くことが期待され、今後のナノカーボン科学を一新する分子です。今回得られたカーボンナノベルトは赤色蛍光を発する有機分子で、発光材料や半導体材料として各種電子デバイスに搭載できる可能性があります。さらに、カーボンナノベルトをテンプレートにした製法で単一構造のカーボンナノチューブが得られれば、軽くて曲げられるディスプレイや省電力の超集積 CPU、バッテリーや太陽電池の効率化など、非常に幅広い応用が期待されます。



図 1. (6,6)Carbon Nanobelt (1)





図 2. (6,6)Carbon Nanotube の模式図



Scheme 1.


文献

- 1) Molecular orbitals in homogenous Reihen mehrkerniger aromatischer Kohlenwasserstoffe: I. Die Eigenwerte von LCAO-MO's in homogenous Reihen
E. Heilbronner, *Helv. Chim. Acta.* **1954**, *37*, 921.
- 2) for example:
a) F. H. Kohnke, A. M. Z. Slawin, J. F. Stoddart, D. J. Williams, *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 892. b) S. Kammermeier, P. G. Jones, R. Herges, *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 2669. c) F. Vögtle, A. Schroder, D. Karbach, *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 575. d) L. T. Scott, *Angew. Chem. Int. Ed.* **2003**, *42*, 4133. e) B. L. Merner, L. N. Dawe, G. J. Bodwell, *Angew. Chem. Int. Ed.* **2009**, *48*, 5487.
- 3) Helical microtubules of graphitic carbon
S. Iijima, *Nature* **1991**, *354*, 56.
- 4) Synthesis of a Carbon Nanobelt
G. Povie, Y. Segawa, T. Nishihara, Y. Miyauchi, K. Itami, *Science* **2017**, *356*, 172.

高性能な液晶性半導体材料：Ph-BTBT-10

D5491 Ph-BTBT-10 (1)

100mg 32,500 円 250mg 68,500 円



Ph-BTBT-10 (1)

液晶性半導体材料 Ph-BTBT-10 (1) は、近年報告された有機半導体材料の中でも特に優れたキャリア輸送特性を示す p 型半導体材料です (図 1)。半那らはスピコート法で作製した FET 素子において、酸化物半導体 (IGZO) と同等の移動度 ($\mu_{\max} = 14.7 \text{ cm}^2/\text{Vs}$) と大気安定性を有することを報告しています¹⁾。そこで、このような特性をもつ液晶性半導体材料である 1 について、蒸着法による OFET 素子の作製および評価検討を行いました。また、1 の蒸着膜の構造解析とアニール処理後の結晶構造の変化について 2D-GIXD による調査・解析も実施しています^{2,3)}。



図 1. OFET 素子構造




図 2. 蒸着型 OFET 素子の伝達特性

(a) w/o annealing (ODTS) (b) annealing 120 °C, 5 min (ODTS)

表 1. Ph-BTBT-10 を用いた蒸着型 OFET 素子特性

Compound	SAM	Annealing Temp. (°C)	Mobility (cm ² /Vs)	V _{th} (V)
Ph-BTBT-10 (1)	bare	w/o	0.87 ~ 0.91	-24
		120	4.24 ~ 4.86	-8
	ODTS	w/o	1.40 ~ 1.42	-23
		120	10.3 ~ 14.0	-22

蒸着法によりトップコンタクト型の FET 素子を作製しました (図 1)。素子の FET 特性を図 2 および表 1 に示します。全ての素子において安定な p 型半導体特性が得られ、かつ、Self-assembled monolayer (SAM) の有無に関わらず、アニーリング処理 (120 °C, 5 min) を行うことで FET 性能の大幅な向上が見られました。bare 基板における素子性能はホール移動度 $\mu_{\max} = 4.86 \text{ cm}^2/\text{Vs}$ 、 $V_{\text{th}} = -8 \text{ V}$ の良好な値が得られ、ODTS 処理基板では閾値電圧は大きい ($V_{\text{th}} = -22 \text{ V}$) ものの I_{DS} が大幅に増加し、ホール移動度 $\mu_{\max} = 14.0 \text{ cm}^2/\text{Vs}$ の良好な値が得られました。




図 3. 2D-GIXD 解析 (加熱 *in-situ* 測定)




図 4. 2D-GIXD 解析 (Out-of-plane, bare 基板)




図 5.1 の相転移イメージ

Ph-BTBT-10 薄膜の加熱 *in situ* 2D-GIXD 測定データを図 3, 4 に示します^{2,3)}。室温および 60 °C にて単分子層に起因するピークが得られ、その層間距離 (d) は 27 Å 程度であることがわかりました。一方、120 °C まで昇温した場合、回折ピークに明確な変化が現れることから、単分子層 ($d = 27 \text{ \AA}$) から 2 分子層 ($d = 54 \text{ \AA}$) への転移が起こったと考えられます (図 4, 5)。液晶相 (SmE) への相転移温度は 144 °C であり¹⁾、180 °C の加熱条件下では SmE に由来する回折ピークが観測されています (図 4)。また、180 °C (SmE) から室温へ急冷却すると、単分子層と 2 分子層の混合層を発現することが示唆され、本系において冷却速度は均一な結晶膜を得るために極めて重要なファクターであると推測されます。

以上の結果から、蒸着法により成膜した Ph-BTBT-10 膜においても、単分子層から 2 分子層へ転移することがわかりました。本解析結果は、実際の FET 特性の大幅な変動と一致します。1 は既に報告されている塗布法に加えて、蒸着法でも使用可能な高い実用的有機半導体材料であることが見出されました。

文献


- 1) Liquid crystals for organic thin-film transistors
H. Iino, T. Usui, J. Hanna, *Nat. Commun.* **2015**, 6, 6828.
- 2) 2D-GIXD experiments were performed at the BL46XU and BL19B2 of SPring-8 with the approval of the Japan Synchrotron Radiation Research Institute (JASRI) (Proposal No. 2017B1817 and 2017B1629).
We acknowledge Prof. Noriyuki Yoshimoto, Assist. Prof. Daiki Kuzuhara and Mr. Shimpei Miura (Iwate Univ.), Assist. Prof. Mitsuharu Suzuki (NAIST) and Dr. Tomoyuki Koganezawa (JASRI) for technical support in GIXD.
- 3) 三浦他, 第 65 回応用物理学会春季学術講演会 20a-P7-1.

光学活性なシクロプロパンの合成に有用な不斉ルテニウム触媒

R0196 Ru(II)-(S)-Pheox Catalyst (1)

200mg 6,600 円 1g 23,100 円

Ru(II)-(S)-Pheox 触媒 (1) は、岩佐らにより開発されたキラルフェニルオキサゾリニルルテニウム (II) 錯体であり、光学活性シクロプロパンの合成に用いることができます。1 の存在下、オレフィン類にジアゾ酢酸エステル^{1-6,8)} あるいはジアゾメチルホスホン酸エステル^{7,8)} を作用させると、エナンチオ選択的なシクロプロパン化反応が進行します。本反応は、医薬開発に有用な光学活性シクロプロパンの合成への利用が期待されます。



文献

- 1) A.-M. Abu-Elfotoh, K. Phomkeona, K. Shibatomi, S. Iwasa, *Angew. Chem. Int. Ed.* **2010**, *49*, 8439.
- 2) S. Chanthamath, K. Phomkeona, K. Shibatomi, S. Iwasa, *Chem. Commun.* **2012**, *48*, 7750.
- 3) S. Chanthamath, D. T. Nguyen, K. Shibatomi, S. Iwasa, *Org. Lett.* **2013**, *15*, 772.
- 4) S. Chanthamath, D. S. Takaki, K. Shibatomi, S. Iwasa, *Angew. Chem. Int. Ed.* **2013**, *52*, 5818.
- 5) S. Chanthamath, H. W. Chua, S. Kimura, K. Shibatomi, S. Iwasa, *Org. Lett.* **2014**, *16*, 3408.
- 6) Y. Nakagawa, S. Chanthamath, K. Shibatomi, S. Iwasa, *Org. Lett.* **2015**, *17*, 2792.
- 7) S. Chanthamath, S. Ozaki, K. Shibatomi, S. Iwasa, *Org. Lett.* **2014**, *16*, 3012.
- 8) S. Chanthamath, S. Iwasa, *Acc. Chem. Res.* **2016**, *46*, 2080.

関連製品

D5368 (S)-Pheox

1g 5,500 円 5g 18,600 円

ペルオキシダーゼ用調製済み基質溶液

A3176 AzBTS (Ready-to-use solution) [for ELISA] (1)

100mL 10,000 円

C3384 4-CN (= 4-Chloro-1-naphthol) (Ready-to-use solution) [for Western blotting] (2)

100mL 10,000 円

ペルオキシダーゼは過酸化水素の存在下、基質を酸化させる作用をもつ酵素で、生化学の実験に広く用いられています。1 および 2 は、ペルオキシダーゼ検出に使用する調製済み基質溶液であり、それぞれ以下に示した手順により検出を行うことができます。

AzBTS (Ready-to-use solution) (1) は基質である 2,2'-アジノビス(3-エチルベンゾチアゾリン-6-スルホン酸 アンモニウム塩) (AzBTS) と過酸化水素を含む ELISA 用調製済み溶液で、ペルオキシダーゼと反応すると可溶性の緑色生成物を生じます。生成物は 405 nm の吸光度で測定可能です。

使用方法：96 ウェルプレートを用いた使用方法を示します。

- 1 を室温に戻し、軽く混ぜる。
- 1 を各ウェルに 100 μ L ずつ加える。
- 室温で 30 分間反応させる。
- 反応開始後、1 時間以内に 405 nm の吸光度を測定する。




図 1. 96 ウェルプレートを用いた使用例

4-CN (Ready-to-use solution) (2) は基質である 4-CN (4-クロロ-1-ナフトール) と過酸化水素を含むウェスタンブロット用調製済み溶液で、ペルオキシダーゼ (HRP) と反応すると沈殿性の青色生成物を生じます。

使用方法：

- HRP 標識された検出用抗体でメンブレン処理し洗浄する。
- メンブレンに 2 を加え発色させる。
- 溶液を捨て、メンブレンに純水を加えて反応を停止させる。




図 2. ウェスタンブロットの例

本製品はいずれも試薬であり、試験・研究用のみにご使用ください。


文献

- 1) *Antibodies, A Laboratory Manual*, ed. by E. Harlow and D. Lane, Cold Spring Harbor Laboratory Press, New York, 1988, p. 404, p. 506, p. 594.

c-Jun N 末端キナーゼ (JNK) 阻害剤 : SU 3327

A2940 SU 3327 (1)

20mg 20,500 円 100mg 71,000 円



SU 3327 (1) は構造活性相関の研究に基づいて開発された JNK (c-Jun N-terminal kinase) の阻害剤です¹⁾。1 は活性化転写因子 -2 (ATF-2) の JNK 1 によるリン酸化を IC₅₀ = 0.7 μM で基質競合的に阻害します。また、1 は JNK と JIP1* のタンパク質間相互作用を IC₅₀ = 239 nM で阻害します。

JNK の一つの機能として、この酵素は Table 1 に示すような刺激によるオートファジーの誘導に関与していることが知られています。マウスのモデル系において急性アルコール処理によるオートファジーの減少が 1 によって防止されることが示されています²⁾。

*JIP1: JNK 相互作用タンパク質 (JNK-interacting protein; JIP) は JNK の足場タンパク質として同定されました^{3,4)}。JIP1 は JIP ファミリーの 1 つで、JNK 結合ドメイン (JNK binding domain; JBD), SRC 相同 (SH3) ドメイン、ホスホチロシン結合 (phosphotyrosine-binding; PTB) ドメインを有しています。JBD はコンセンサス配列 R/KXXXXLXL を有します⁵⁾。文献 1 で使用されているプローブ pep-JIP1 にはこの配列が含まれます。

Table 1. Autophagy by various stimuli implicating JNK

Stimulation	References
Neuronal excitotoxic stimuli	<i>Eur. J. Neurosci.</i> 2003 , <i>18</i> , 473.
Caspase inhibition	<i>Science</i> 2004 , <i>304</i> , 1500.
Starvation	<i>J. Immunol.</i> 2006 , <i>177</i> , 5163.
T-cell receptor activation	<i>J. Immunol.</i> 2006 , <i>177</i> , 5163.
Cytokine stimulation	<i>Immunol. Cell Biol.</i> 2006 , <i>84</i> , 448.
ER stress	<i>Mol. Cell. Biol.</i> 2006 , <i>26</i> , 9220.

本製品は試薬であり、試験・研究用のみにご使用ください。


文献

- Design, synthesis, and structure-activity relationship of substrate competitive, selective, and *in vivo* active triazole and thiazole inhibitors of the c-Jun N-terminal kinase
S. K. De, J. L. Stebbins, L.-H. Chen, M. Riel-Mehan, T. Machleidt, R. Dahl, H. Yuan, A. Emdadi, E. Barile, V. Chen, R. Murphy, M. Pellecchia, *J. Med. Chem.* **2009**, *52*, 1943.
- Cytochrome P450E1, oxidative stress, JNK and autophagy in acute alcohol-induced fatty liver
L. Yang, D.-F. Wu, X.-D. Wang, A. I. Cederbaum, *Free Radic. Biol. Med.* **2012**, *53*, 1170.
- Scaffold proteins of MAP-kinase modules
D. N. Dhanasekaran, K. Kashef, C. M. Lee, H. Xu, E. P. Reddy, *Oncogene* **2007**, *26*, 3185.
- A mammalian Scaffold complex that selectively mediated MAP kinase activation
A. J. Whitmarsh, J. Cavanagh, C. Tournier, J. Yasuda, R. J. Davis, *Science* **1998**, *281*, 1671.
- Differential targeting of MAP kinases to the ETS-domain transcription factor Elk-1
S.-H. Yang, A. J. Whitmarsh, R. J. Davis, A. D. Sharrocks, *EMBO J.* **1998**, *17*, 1740.

シクロオキシゲナーゼ (COX) 阻害剤 : フルルビプロフェン

F0371 Flurbiprofen (1)

5g 10,400円 25g 30,700円



フルルビプロフェン (1) はシクロオキシゲナーゼ (COX) 阻害剤の一種で, COX-1 と COX-2 をそれぞれ 0.1 μM , 0.4 μM の IC_{50} で阻害します¹⁾。1 はラセミ体混合物として存在し, S 体は R 体よりも強く COX イソ酵素を阻害します^{2,3)}。また, 1 は非ステロイド性抗炎症剤として広く知られています¹⁾。さらに 1 は, ヒト胚性幹細胞の自己複製と多機能性を維持します^{4,5)}。


* 文献 1 では組み替え体のヒト COX-1 と COX-2 が使用されています。

本製品は試薬であり, 試験・研究用のみにご使用ください。

文献

- 1) Expression and selective inhibition of the constitutive and inducible forms of human cyclo-oxygenase
J. K. Gierse, S. D. Hauser, D. P. Creely, C. Koboldt, S. H. Rangwala, P. C. Isakson, K. Seibert, *Biochem. J.* **1995**, 305, 479.
- 2) Stoichiometric and kinetics of the interaction of prostaglandin H synthase with anti-inflammatory agents
R. J. Kulmacz, W. E. Lands, *J. Biol. Chem.* **1985**, 260, 12572.
- 3) Flurprofen and enantiomers in ophthalmic solution tested as inhibitors of prostanoid synthesis in human blood
N. J. van Haeringen, A. A. van Sorge, J. L. van Delft, V. M. W. Carballosa Coré-Bodelier, *J. Ocul. Pharmacol. Ther.* **2000**, 16, 345.
- 4) High-throughput screening assay for the identification of compounds regulating self-renewal and differentiation in human embryonic stem cells
S. C. Desbordes, D. G. Placantonakis, A. Ciro, N. D. Socci, G. Lee, H. Djaballah, L. Studer, *Cell Stem Cell* **2008**, 2, 602.
- 5) Histone modification profiling reveals differential signatures associated with human embryonic stem cell self-renewal and differentiation
N. V. Bhanu, S. Sidoli, B. A. Garcia, *Proteomics* **2016**, 16, 448.

糖鎖関連試薬カタログ <第5版・2017年7月発行>



Reagents for
Glyco Chemistry & Biology
5th Edition

← 新製品200品目追加 (全900品目収録)

- N結合型糖鎖(合成品)を新規収録
- 糖鎖コンジュゲートに応用可能な糖鎖を多数追加
- 約250品目の ¹H-NMRスペクトルデータを収録
- 糖の種類ごとに収録し、わかりやすく紹介

糖鎖研究に役立つ各種試薬を多数収録

ホームページでのご請求は ▶▶▶ www.TCIchemicals.com/ja/jp/tosa-catalog-tm/

出展のご案内
ぜひお立ち寄りください



第37回日本糖質学会年会

2018年8月28日(火)~30日(木) 仙台国際センター

第48回複素環化学討論会

2018年9月3日(月)~5日(水) 長崎ブリックホール

第91回日本生化学会大会

2018年9月24日(月)~26日(水) 国立京都国際会館

第77回日本癌学会学術総会

2018年9月27日(木)~29日(土) 大阪国際会議場(グランキューブ大阪)

オンラインカタログ

構造式、品名(和・英)、分子式、CAS番号、キーワード、
弊社製品コードからの検索が可能です。

www.TCIchemicals.com/ja/jp/



ご注文・カタログの
ご請求は

最寄りの弊社製品取扱店へ

お問い合わせは

○ご注文・カタログのご請求に関して
東京化成販売(株) Tel:03-3668-0489 Fax:03-3668-0520
大阪営業部 Tel:06-6228-1155 Fax:06-6228-1158

○製品に関して
學術部 Tel:03-5640-8857 Fax:03-5640-8868
E-mail: information@TCIchemicals.com

TCI 東京化成工業株式会社

〒103-0023 東京都中央区日本橋本町4-10-2 www.TCIchemicals.com/ja/jp/

本文に掲載した化学品は試薬であり、試験・研究用のみに使用するものです。化学知識のある専門家以外の方の使用はお避けください。
弊社は掲載した製品に関して発生した特許法上の諸問題をユーザーの方々に保証するものではありません。
掲載した製品およびその価格等は発行時のものです。諸事情によりやむを得ず変更を行う場合があります。
本誌の内容の一部または全部を無断で転載あるいは複製することはご遠慮ください。

Printed in Japan